

HERPES, ORAL

Commonly, these infections are acquired for the first time in early childhood and may reappear throughout a person's lifetime (recurrent infection).

CAUSE	Herpes simplex virus type 1 (HSV-1).
SYMPTOMS	Primary infection causes blister-like sores inside the mouth (gingivostomatitis) and on the gums. Fever and irritability may also occur. Recurrent infections (cold sores, fever blisters) appear on lips and face. Some children may have the virus in their bodies, but do not develop symptoms. Herpes simplex virus can also cause infections of the eyes, fingers, and central nervous system.
SPREAD	By close person-to-person contact, such as through direct contact with saliva (e.g., kissing) or touching the fluid from the blisters or sores. Most experts believe that herpes is not spread from lipsticks, towels, washcloths, drinking glasses, or toys. However, to prevent spread of other infectious germs, personal items should not be shared.
INCUBATION	It takes 2 to 14 days from the time a person is exposed for the first time until symptoms occur.
CONTAGIOUS PERIOD	First occurrence (primary infection): up to 2 weeks, occasionally up to 7 weeks. Recurrent infection: usually 3 to 5 days.
EXCLUSION	<u>Childcare</u> : Primary infection - Until those children who do not have control of their oral secretions no longer have active sores inside the mouth. Recurrent infections (fever blisters and cold sores) - None. <u>School</u> : None.
TREATMENT	A topical medication may be prescribed.

PREVENTION/CONTROL

- Encourage children not to touch the sores.
- Wash hands thoroughly with soap and warm running water after having contact with the sores or saliva. **Thorough handwashing is the best way to prevent the spread of communicable diseases.**
- Wear disposable medical gloves when touching the sores is necessary (e.g., when applying medication).
- DO NOT kiss an infected person on or near the mouth when sores are present.
- Clean and sanitize mouthed toys, objects, and surfaces at least daily and when soiled. (See pgs 34-36.)

For more information, call Hennepin County HSPHD-Epidemiology at (612) 543-5230 or call your local health department.

Prepared by Hennepin County Human Services and Public Health Department (HSPHD)

June 2008